

## *In vitro* and *in vivo* assessment of free radical scavenging and antioxidant activities of *Veronica persica* Poir

Javad Sharifi-Rad<sup>1</sup>, Mehdi Sharifi-Rad<sup>2\*</sup>, Bahare Salehi<sup>3,4\*</sup>, Marcello Iriti<sup>5</sup>, Amir Roointan<sup>6</sup>, Dima Mnayer<sup>7</sup>, Azam Soltani-Nejad<sup>8</sup>, Azam Afshari<sup>9</sup>

<sup>1</sup>Phytochemistry Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>2</sup>Department of Medical Parasitology, Zabol University of Medical Sciences, 61663335 Zabol, Iran

<sup>3</sup>Medical Ethics and Law Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>4</sup>Student Research Committee, Shahid Beheshti University of Medical Sciences, 22439789 Tehran, Iran

<sup>5</sup>Department of Agricultural and Environmental Sciences, Milan State University, 20133 Milan, Italy

<sup>6</sup>Department of Medical Biotechnology, School of Medicine, Fasa University of Medical Sciences, Fasa, Iran

<sup>7</sup>Faculty of Sciences, Lebanese University, Beirut 6573, Lebanon

<sup>8</sup>Department of Genetics and Biotechnology, Osmania University, Hyderabad, India

<sup>9</sup>Department Rangeland and Watershed, Faculty of Water and Soil, University of Zabol, Zabol, Iran

**Correspondence to:** mehdi\_sharifrad@yahoo.com; bahar.salehi007@gmail.com

**Received** December 15, 2017; **Accepted** January 29, 2018; **Published** June 25, 2018

**Doi:** <http://dx.doi.org/10.14715/cmb/2018.64.8.9>

**Copyright:** © 2018 by the C.M.B. Association. All rights reserved.

**Abstract:** With the appearance of new disorders along with inability of some conventional therapies for the treatment of diseases without any side effects, the discovery of safe and efficient therapeutic agents is of utmost importance in the medical area. In this context, medicinal plants as promising therapeutic candidates can provide a reliable and efficient profile. Since free radicals are at the center of various disorder pathways, reducing their production or complete removal of these chemical species could be advantageous for prevention and treatment of many diseases. In this experiment, free radical scavenging and antioxidant activities of *Veronica persica* Poir., a known medicinal plant, were evaluated using *in vitro* and *in vivo* assays. Chemical characterization results showed a high phenolic content in the *V. persica* methanol extract. In addition, *in vitro* assays including DPPH radical-scavenging assay, nitric oxide-scavenging activity assay, hydrogen peroxide scavenging test and bleomycin-dependent DNA damage test revealed significant antioxidant power and radical scavenging capacity of this plant. In accordance, *in vivo* experiments showed inhibitory effects of the methanol extract on lipid peroxidation, a main cause of cell damage. Our findings revealed the promising potential of this plant in reducing free radicals through different pathways. Moreover, our data suggested a correlation between the high phenolic content of the *V. persica* extract and its free radical scavenging and antioxidant activities.

**Key words:** Plantaginaceae; Methanol extract; Lipid peroxidation; Oxidative stress; Polyphenols; ROS.

### Introduction

Free radicals are chemical species that contain one or more unpaired electrons, which can be either neutral or positively or negatively charged. The important ones in the human body are superoxide anion ( $O_2^-$ ), hydroxyl radical ( $OH^\cdot$ ) and nitric oxide ( $NO^\cdot$ ), with the ability to produce many other free radicals mostly derived from unsaturated fatty acids (1). These molecules with unstable nature have the affinity to reach stability by donating their unpaired electron to other molecules or taking another electron from other cell molecules (1). In low to moderate concentrations, these highly reactive fragmented atoms can play essential roles in hormonal regulations, signal transduction pathways, acting as transcription factors, upregulating defense genes and adaptive responses. However, they can be harmful in higher concentrations (2). They also play a critical role in the pathogenesis of various chronic degenerative disorders, including atherosclerosis, cancer, neurodegenerative diseases, diabetes mellitus, aging, rheumatoid arthritis and inflammation (3-5). Therefore, reactive oxygen

species (ROS) and reactive nitrogen species (RNS) play a dual role. ROS and RNS overproduction causes oxidative and nitrosative stress which leads to impairment of cell macromolecules, including lipids, DNA and proteins. However, moderate concentrations of ROS and RNS are beneficial during infections (the respiratory burst of macrophages), in cell signaling and mitogenic stimulation (2, 6). This delicate balance is accomplished by a mechanism called “redox regulation” (2). Synthetic antioxidants in foods might be a suitable choice; however, using such additives can result in toxicity since they contain phenolic compounds with various degrees of alkyl substitution. On the other hand, natural antioxidants include phenolic compounds, nitrogen-containing compounds, carotenoids, tocopherols, ascorbic acid and their derivatives (7, 8). These natural antioxidants possess redox properties giving them the ability to act as reducing agents or hydrogen-atom donors (9). Therefore, they can function as free-radical scavengers and chain breakers, chelators of pro-oxidant metal ions and quenchers of singlet-oxygen formation (10).

Plant-based foods can reduce ROS activities through

different mechanisms including *i*) avoiding ROS generation; *ii*) scavenging ROS and their byproducts; *iii*) exchanging ROS with less reactive molecules, or *iv*) improving the resistance of sensitive biological targets to ROS damages (11).

Until now, various investigations have been conducted to find any natural agent able to reduce free radical damages (12, 13). In the search for naturally occurring antioxidants from plant sources, *Veronica* species were found to have significant radical scavenging activity (11). The genus *Veronica* (Plantaginaceae) includes annual and perennial plants located in central and southern Europe and Turkey, and is represented by 79 species, 26 of which are endemic in the Turkish flora (12). Based on some investigations, these species have been traditionally used in the treatment of various diseases such as cancer, influenza, hemoptysis, laryngopharyngitis, hernia and respiratory disorders (14, 15). The genus *Veronica* is believed to contain mostly iridoid glucosides, especially benzoic and cinnamic acid esters of catalpol, as well as some phenylethanoid and flavonoid glycosides (4). These flavonoids were found to be flavone glycosides often with additional hydroxyl substitution at C-6 or C-8 of the A-ring (12, 16). Iridoid glucosides, phenylethanoid glycosides, and alkaloids have also been reported from *Veronica persica* Poir. in previous studies (4). Therefore, our study aims to check free radical scavenging and antioxidant activities of *V. persica* using different *in vitro* and *in vivo* assays.

## Materials and Methods

### Plant material and preparation of the extract

Aerial parts (stems, leaves, and flowers) of *Veronica persica* Poir. were collected at flowering stage in April 2016 from wild plants in the mountains of Meymand, Firuzabad County, Fars Province, Iran (Coordinates: 28°52'04"N 52°45'12"E). The plant was taxonomically identified by a botanist. All parts of the plants were dried in the shade, pulverized into a fine powder using a grinder sieved through a No. 22 mesh sieve and stored in an air-tight container until analysis. A volume of 200 mL of 70% methanol was added to 20 g of powder and kept on a mechanical shaker for 72 h. The content was filtered and concentrated under reduced pressure and controlled temperature to obtain a dark gummy residue. The concentrated extract was stored dry in an amber-colored flask at 4°C for future analysis.

### Animals

In this study, all animal manipulations were carried out according to the Helsinki Convention. The protocol study was approved by the local Ethical Committee (IR.SBMU.RETECH.REC.1396.586). Male Wistar rats (190 ± 4.5 g) and used for *in vivo* assays. The animals were kept in clean metabolic stainless-steel cages placed in a well-ventilated room and allowed to adapt to their controlled environmental conditions (24 ± 5 °C and 12:12 hours light-dark cycle) before and during the tests. The animals were having free access to food and water. All animal tests were done during the light period.

## Polyphenol contents

### Total phenolics

The content of total phenolics in the *V. persica* methanol extract was determined spectrophotometrically by the Folin Ciocalteu reagent with the slightly modified method of Wolfe *et al.* (17). An aliquot of the extract (0.5 ml) was mixed with 3 ml of 10% Folin-Ciocalteu reagent and 2.5 ml of 75% w/v sodium carbonate (Sigma-Aldrich, St. Louis, Missouri, United States) solution. The resulting mixture was vortexed for 20 seconds and incubated at 40°C for 25 min for color development. The absorbance of the samples was measured at 765 nm against a blank in a Hitachi U-2001 spectrophotometer (Tokyo, Japan). All measurements were carried in triplicate and the total phenolic content was expressed as mg/g tannic acid equivalent.

### Total flavonoids

The determination of total flavonoid content of the plant extract was performed according to Ordon *et al.* (2006) method (18). Briefly, a volume of 0.5 ml of 2% AlCl<sub>3</sub> ethanol solution was added to 0.5 mL of the extract. After one hour of incubation at room temperature (25 ± 3 °C), the absorbance was measured at 420 nm against a blank in a Hitachi U-2001 spectrophotometer (Tokyo, Japan). All measurements were carried in triplicate and the total flavonoid content was expressed as mg/g quercetin equivalent.

### Total flavonols

The determination of total flavonol contents of the plant extract was performed according to the method described by Kumaran and Karunakaran (2007) (19). The reacting mixture consisted of 3 mL of the sample, 3 ml of AlCl<sub>3</sub> prepared in ethanol and 3.0 ml of (50 g/L) sodium acetate solution. The absorption at 440 nm was measured against a blank after 3 h at 20°C by using Hitachi U-2001 spectrophotometer (Tokyo, Japan). All measurements were carried in triplicate and the total flavonol content was calculated as quercetin equivalent (mg/g).

### Total proanthocyanidins

The determination of total proanthocyanidins was performed according to Sun *et al.* (1998) (20) with slight modifications. Briefly, 0.5 ml of 1 mg/mL extract solution was added 3 ml of vanillin-methanol (4 % v/v) and 1.5 ml of hydrochloric acid and, then, the mixture was vortexed shortly. The absorbance of the mixture was measured at 500 nm against a blank after 15 min at room temperature (25 ± 5 °C). All measurements were carried in triplicate and the total proanthocyanidin content was expressed as catechin equivalents (mg/g).

## In vitro antioxidant activities

### DPPH radical-scavenging assay

The scavenging capacity of the plant extract against the stable free radical 2,2-diphenyl-1-picrylhydrazyl (DPPH, Sigma-Aldrich, St. Louis, MO, USA) was measured according to the method of Liyana-Pathiranan and Shahidi (2005) (21). Briefly, a solution of 0.135 mM DPPH in methanol was prepared, and 1.0 ml of this solution was mixed with 1.0 mL of the plant extract prepa-

red in methanol ranging from 0.025 to 0.5 mg. Butylated hydroxytoluene (BHT) and rutin were used as standard drugs separately. The reaction mixture was vortexed thoroughly and left in the dark at room temperature ( $25 \pm 5^\circ\text{C}$ ) for 30 min. The absorbance of the mixture was measured at 517 nm against a blank by using Hitachi U-2001 spectrophotometer (Tokyo, Japan). The ability of the plant extract to scavenge DPPH radical was calculated by the equation:

$$\text{DPPH radical scavenging activity} = \frac{(\text{Absorbance}_{\text{control}} - \text{Absorbance}_{\text{sample}})}{(\text{Absorbance}_{\text{control}})} \times 100$$

Where: absorbance control is the absorbance of DPPH radical plus methanol; absorbance sample is the absorbance of DPPH radical plus sample extract or standard.

### Nitric oxide-scavenging activity

For the determination of nitric oxide (NO) scavenging activity of the plant extract the method of Garrat (1964) was used (22). Briefly, 2 mL of 10 mM sodium nitroprusside prepared in 0.5 mL phosphate buffer saline (pH 7.4) were mixed with 0.5 mL of the plant extract. Butylated hydroxytoluene (BHT) and rutin were used at different concentrations (0.025 - 0.5 mg/mL). The mixture was incubated at  $25^\circ\text{C}$ . After 3 h, 0.5 mL of incubation solution was withdrawn and mixed with 0.5 mL of Griess reagent [1.0 mL sulfanilic acid reagent (0.33 % prepared in 20 % glacial acetic acid) at room temperature ( $25 \pm 5^\circ\text{C}$ ) for 5 min with 1 mL of naphthylethylenediamine dichloride (0.1% w/v)]. The mixture was incubated at room temperature ( $25 \pm 5^\circ\text{C}$ ) for 30 min, followed by the measurement of absorbance at 540 nm against a blank using Hitachi U-2001 spectrophotometer (Tokyo, Japan). The amount of nitric oxide radical inhibited by the extract was calculated using the following formula:

$$\text{NO radical scavenging activity} = \frac{(\text{Absorbance}_{\text{control}} - \text{Absorbance}_{\text{sample}})}{(\text{Absorbance}_{\text{control}})} \times 100$$

Where: absorbance control is the absorbance of NO radical plus methanol; absorbance sample is the absorbance of NO radical plus sample extract or standard.

### Hydrogen peroxide scavenging test

The method of Ruch *et al.* (1989) was used to estimate the hydrogen peroxide scavenging activity of the plant extract (23). Briefly, 4 mL of the plant extract prepared in distilled water at different concentrations were mixed with 0.6 mL of 4 mM hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) solution prepared in phosphate buffer (0.1 M, pH 7.4) and incubated for 15 min. The absorbance of the solution was read at 230 nm against blank solution (plant extract without  $\text{H}_2\text{O}_2$ ) by using Hitachi U-2001 spectrophotometer (Tokyo, Japan). The amount of hydrogen peroxide radical inhibited by the extract was calculated using the following equation:

$$\text{H}_2\text{O}_2 \text{ radical scavenging activity} = \frac{(\text{Absorbance}_{\text{control}} - \text{Absorbance}_{\text{sample}})}{(\text{Absorbance}_{\text{control}})} \times 100$$

Where: absorbance control is the absorbance of  $\text{H}_2\text{O}_2$  radical plus methanol; absorbance sample is the absorbance of  $\text{H}_2\text{O}_2$  radical plus sample extract or standard.

### Bleomycin-dependent DNA damage

In this test, the pro-oxidant activity of the plant extract and the standard (ascorbic acid) were evaluated by the Ng *et al.*, (2003) method (24). Briefly, the reaction

mixture contained 50  $\mu\text{g}$  of 1 mL bleomycin sulfate, 1 mL of 5 mM magnesium chloride, 1 mL of 50  $\mu\text{M}$  ferric chloride, 0.5 mL calf thymus DNA (10  $\mu\text{g}/\text{mL}$ ), and 1 mL of various concentrations of the plant extract. The mixture was incubated at  $37^\circ\text{C}$  for 1 h. The reaction was terminated by addition of 0.05 mL ethylenediaminetetraacetic acid (EDTA) (0.1 M). The color was developed by adding 0.5 mL thiobarbituric acid (TBA) (1% w/v) and 0.5 mL hydrochloric acid (25% v/v) followed by incubation at  $37^\circ\text{C}$  for 20 min. After centrifugation, the extent of DNA damage was measured at 532 nm by using Hitachi U-2001 spectrophotometer (Tokyo, Japan). The ascorbic acid was the positive control.

### Calculation of 50% inhibitory concentration ( $\text{IC}_{50}$ )

The concentration (mg/mL) of the extract required to scavenge 50% of the radicals was calculated by using the percentage scavenging activities at various different concentrations of the extract. The  $\text{IC}_{50}$  was calculated using the formula (25):

$$\text{EXP}(\text{LN}(\text{conc} > 50\%) - ((\text{signal} > 50\% - 50) / (\text{signal} > 50\% - \text{signal} < 50\%)) * \text{LN}(\text{conc} > 50\% / \text{conc} < 50\%))$$

### In vivo study

#### Animal grouping and extract administration

Thirty-five male Wistar rats were randomized into five groups consisting of 7 each. The first group served as control and was given 0.5 mL distilled water alone per day for one week with the aid of oropharyngeal cannula. In the second group, animals served as hepatotoxic control, treated with carbon tetrachloride ( $\text{CCl}_4$ ) in a single dose of 0.5 mL administered orally for one week. Groups 3 - 5 were treated like the hepatotoxic control except that they received 0.5 mL of the extract corresponding to 250, 500 and 1000 mg/kg body weight respectively. All the Wistar rats from each group were sacrificed by ether anesthesia. The liver from each animal was excised, rinsed in ice cold 0.25 M sucrose solution and 10% w/v homogenate was prepared in 0.05 M phosphate buffer (pH 7) and centrifuged at  $12,000 \times g$  for 60 min at  $4^\circ\text{C}$ . The supernatant obtained was used for the estimation of superoxide dismutase and catalase activities, lipid peroxidation and reduced glutathione.

#### Superoxide dismutase activity

Superoxide dismutase (SOD) activity was evaluated with the method described by Misra and Fridovich (1972) (26). Briefly, a mixture was prepared contained 0.5 mL of hepatic post-mitochondrial supernatant (PMS), 0.4 mL of 25  $\mu\text{M}$  nitroblue tetrazolium, 1 mL of 50 mM sodium carbonate, and 0.2 mL of freshly prepared 0.1 mM hydroxylaminehydrochloride. The reaction mixture was mixed quickly by inversion followed by the addition of supernatant of 0.1 mL of liver homogenate (10% w/v). The change in absorbance was read at 560 nm by using Hitachi U-2001 spectrophotometer (Tokyo, Japan). Percentage inhibition was calculated using this equation:

$$\% \text{ Inhibition of superoxide dismutase} = \frac{(\text{Normal activity} - \text{Inhibited activity})}{(\text{Normal activity})} \times 100$$

#### Lipid peroxidation assay

Lipid peroxidation in the liver was measured using thiobarbituric acid reactive substances (TBARS) accor-

ding to the slightly modified method of Niehius and Samuelsson (1968) (27) Briefly, 0.1 mL of liver homogenate (10% w/v) was treated with 2 mL of (1:1:1 ratio) thiobarbituric acid (TBA)-trichloroacetic acid (TCA)-HCl reagent (TBA 0.37%, TCA 15% and 0.25 N HCl). All the tubes were placed in a boiling water bath for 30 min and then cooled. The amount of malondialdehyde formed in each of the samples was evaluated by measuring the absorbance of the clear supernatant at 535 nm against blank by using Hitachi U-2001 spectrophotometer (Tokyo, Japan). Percentage inhibition was calculated using the equation:

% Inhibition of lipid peroxidation =  $(A_c - A_s)/A_c \times 100$   
 Where:  $A_c$  is the absorbance of the control and  $A_s$  is the absorbance of the sample extract.

**Catalase activity assay**

The activity of catalase (CAT) was assayed by a method described by Pari and Latha (2004) (28). In brief, the liver was homogenized in 0.01 M phosphate buffer (pH 7.0) and centrifuge at 5000 rpm. The reaction mixture consisted of 1 mL of 0.01 M phosphate buffer (pH 7.0), 0.4 mL of hydrogen peroxide (0.2 M), and 0.1 mL of liver homogenate (10% w/v). The reaction of the mixture was stopped by adding 2 mL of dichromate-acetic acid reagent (5%  $K_2Cr_2O_7$  prepared with glacial acetic acid). The changes in the absorbance were measured at 620 nm and recorded by using Hitachi U-2001 spectrophotometer (Tokyo, Japan). Percentage of inhibition was calculated using the equation:

% Inhibition of catalase =  $[(\text{Normal activity} - \text{Inhibited activity}) / (\text{Normal activity})] \times 100$

**Reduced glutathione assay**

To determine the reduced glutathione (GSH), the Ellman (1951) method (29) with slight modifications was used. Briefly, an aliquot of 1 mL of the supernatant of liver homogenate was treated with 0.5 mL of Ellman’s reagent (19.8 mg of 5,5’-dithiobisnitro benzoic acid (DTNB) in 100 mL of 0.1% sodium nitrate) and 3 mL of phosphate buffer (0.2 M, pH 8.0). The absorbance was measured at 412 nm by using Hitachi U-2001 spectrophotometer (Tokyo, Japan). The percentage inhibition of reduced glutathione was calculated using the following equation:


% Inhibition of reduced glutathione =  $(A_c - A_s)/A_c \times 100$   
 Where:  $A_c$  is the absorbance of the control and  $A_s$  is the absorbance of the sample extract.

**Statistical analysis**


The experimental results were expressed as the mean  $\pm$  standard deviation (SD) of three replicates and were subjected to paired student’s t-test. Significant levels were tested at  $P < 0.05$ .

**Results**

After determination of polyphenol contents in the *V. persica* extract, a high total phenol content (88.5 mg/g tannic acid equivalent). The amounts of total flavonoids, proanthocyanidins and flavonols in the plant extract were 8.3 mg/g quercetin equivalent, 7.1 mg/g catechin equivalent and 6.5 mg/g quercetin equivalent, respectively (Figure 1).



**Figure 1.** Polyphenol contents of *V. persica* extract.



**Figure 2.** DPPH radical scavenging activity of *V. persica* extract

The *V. persica* extract exhibited a dose-dependent *in vitro* DPPH scavenging capacity ( $IC_{50}$  0.03 mg/mL) (Figure 2). This antiradical activity was the weakest among the other reactive species evaluated.

The NO scavenging power test showed the ability of *V. persica* to inhibit NO in a moderate dose-dependent manner with an  $IC_{50}$  of 0.09 mg/mL, as shown in table 1.

The antioxidant activity of the plant extract was also assessed using hydrogen peroxide scavenging test. *V. persica* extract exhibited a dose-dependent scavenging capacity against  $H_2O_2$  when compared with vitamin C and BHT used as controls (Table 2). The  $IC_{50}$  value for

**Table 1.** Nitric oxide radical scavenging activity of *V. persica* extract.

Sample	Concentration (mg/mL)	% Inhibition	$IC_{50}$
Plant extract	0.025	15.42 $\pm$ 0.29	0.09
	0.05	45.22 $\pm$ 0.01	
	0.1	51.11 $\pm$ 0.04	
	0.2	54.79 $\pm$ 0.65	
	0.5	59.80 $\pm$ 0.11	
BHT			< 0.025
Ascorbic acid			< 0.025

Values are expressed as mean  $\pm$  SD.

**Table 2.** Hydrogen peroxide radical scavenging activity of *V. persica* extract.

Sample	Concentration (mg/mL)	% Inhibition	$IC_{50}$
Plant extract	0.025	35.29 $\pm$ 0.11	0.04
	0.05	54.17 $\pm$ 0.00	
	0.1	65.62 $\pm$ 0.41	
	0.2	72.44 $\pm$ 0.1	
	0.5	80.80 $\pm$ 0.09	
BHT			< 0.025
Ascorbic acid			< 0.025

Values are expressed as mean  $\pm$  SD.

**Table 3.** Pro-oxidant activity of *V. persica* extract.

Sample	Concentration ( $\mu\text{g/mL}$ )				
	25	50	100	200	400
Plant extract	0.19 $\pm$ 0.04	0.16 $\pm$ 0.01	0.61 $\pm$ 0.00	0.08 $\pm$ 0.02	0.04 $\pm$ 0.01
Ascorbic acid	0.97 $\pm$ 0.04	0.75 $\pm$ 0.05	0.21 $\pm$ 0.09	0.03 $\pm$ 0.01	0.02 $\pm$ 0.01

Values are expressed as mean  $\pm$  SD.

the plant extract was 0.04 mg/ml, whereas vitamin C and BHT showed IC<sub>50</sub> values less than 0.025 mg/ml.

The pro-oxidant activity of the plant extract and the standard (ascorbic acid) was evaluated by their damaging effects to DNA in the presence of the bleomycin-Fe<sup>2+</sup> complex. It was observed that the plant extract is devoid of pro-oxidant activity since the absorbance values decreased with the increase in the extract concentrations Table 3. This result was compared to ascorbic acid effect ( $P < 0.05$ ).

The *in vivo* experiment after administration of different *V. persica* extract amounts (250, 500 and 1000 mg/kg body weight) revealed that the ability of the plant extract in the promotion of CAT and SOD activities, and GSH levels is in a dose-dependent manner ( $P < 0.05$ ) (Table 4). Lipid peroxidation assay in rats treated with the plant extract demonstrated a remarkable increase in thiobarbituric acid reactive substances (TBARS) when compared to normal control group ( $P < 0.05$ ).

## Discussion

Medicinal plants have been applied for the treatment of various disorders since ancient times (30). From a simple cold to a complex disease, such plants can be used as suitable curing agents. Therefore, the discovery of new (phyto) therapeutic agents is of outmost importance in the medical area (31-37). Medicinal plants can be suitable candidates in this context (9, 38-49). *V. persica*, as a well-known medicinal plant, was selected for this study primarily because of its anti-inflammatory potential. Indeed, *V. persica*, also known as birdeye speedwell, a very common annual herb, has a long record in folk medicine (50, 51).

Until now, different investigations have been conducted to find biological activities of *V. persica*. In a study performed by Harput *et al.*, (2002), the activities of five *Veronica* species including *V. cymbalaria*, *V. hederifolia*, *V. pectinata* var. *glandulosa*, *V. persica* and *V. polita* were evaluated (14). The results of their study showed that methanol extracts of all species exhibited inhibitory effects on nitric oxide (NO) production in lipopolysaccharide (LPS)-stimulated macrophages. The extracts also showed cytotoxic activity against KB

epidermoid carcinoma and B16 melanoma cells. Such results indicated their radical scavenging activity which was compared to well-known antioxidants such as BHA and dl- $\alpha$ -tocopherol (14). In another recent experiment, the hydroalcoholic extract of *V. persica* was evaluated with various assays to study the antioxidant, anti-fungal and anti-inflammatory potentials of this plant (51). The chemical composition of the plant extract was characterized by high performance liquid chromatography (HPLC) which quantified 13 compounds. The extract showed a tremendous antioxidant activity using DPPH (79.91% inhibition) and phosphomolybdate (324.4 mg/g ascorbic acid equivalents) assays. The extract also exhibited an antifungal activity against *Aspergillus niger* and *Penicillium hirsutum*. Moreover, the anti-inflammatory activity of *V. persica* extract was confirmed using inflammation experimental models. Anti-inflammatory (52), antioxidant, antibacterial (53, 54), antimutagenic and antitumor (55) activities of different *Veronica* species were also shown by various investigations. These reports are evidences of the enormous potential of *Veronica* species in biomedical sector.

In this experiment, we evaluated the *in vitro* and *in vivo* antioxidant and radical scavenging activities of *V. persica* methanol extract using different assays. The analysis of the polyphenol content showed a low amount of total flavonoids, proanthocyanidins and flavonols in the extract. On the other hand, a high amount of total phenolic was found in the extract. Results of all *in vitro* antioxidant assays including DPPH radical-scavenging, nitric oxide-scavenging and hydrogen peroxide-scavenging tests showed a remarkable antioxidant activity of the *V. persica* extract. DPPH assay results indicated a dose-dependent scavenging ability of the plant extract. However, this ability was not as high as the ascorbic acid or BHA one. The activity of the extract in scavenging NO also showed a dose-dependent trend, though with a higher IC<sub>50</sub>. A similar pattern was observed in H<sub>2</sub>O<sub>2</sub> scavenging test, where the extract exhibited a high capacity against hydrogen peroxide.

In addition, the plant extract did not display pro-oxidant activity. Since the extract contained phenolics, flavonoids, proanthocyanidins and flavonols, the observed antioxidant effects of *V. persica* extract could be

**Table 4.** *V. persica* extract effects on lipid peroxidation, antioxidant enzymes and glutathione in CCl<sub>4</sub> induced liver damage in male Wistar rats.

Groups	Catalase	Superoxide dismutase	Glutathione	Lipid peroxidation
Control	105.42 $\pm$ 0.01 <sup>a</sup>	98.95 $\pm$ 0.11 <sup>a</sup>	90.44 $\pm$ 0.05 <sup>a</sup>	114.85 $\pm$ 0.29 <sup>a</sup>
Carbon tetrachloride	65.58 $\pm$ 0.12 <sup>b</sup>	39.33 $\pm$ 0.01 <sup>b</sup>	29.45 $\pm$ 0.12 <sup>b</sup>	39.34 $\pm$ 0.06 <sup>b</sup>
Carbon tetrachloride + <i>V. persica</i> (C1)	75.33 $\pm$ 0.19 <sup>c</sup>	58.21 $\pm$ 0.09 <sup>c</sup>	55.39 $\pm$ 0.02 <sup>c</sup>	65.92 $\pm$ 0.03 <sup>c</sup>
Carbon tetrachloride + <i>V. persica</i> (C2)	83.47 $\pm$ 0.03 <sup>c</sup>	75.32 $\pm$ 0.14 <sup>d</sup>	62.75 $\pm$ 0.23 <sup>d</sup>	73.29 $\pm$ 0.01 <sup>d</sup>
Carbon tetrachloride + <i>V. persica</i> (C3)	99.89 $\pm$ 0.51 <sup>d</sup>	83.33 $\pm$ 0.09 <sup>e</sup>	69.72 $\pm$ 0.29 <sup>e</sup>	83.72 $\pm$ 0.42 <sup>e</sup>

Values are mean of  $\pm$  SD (n = 7 rats). C1 = 250 mg/kg body wt; C2 = 500 mg/kg body wt; C3 = 1000 mg/kg body wt. Results are expressed as percentage inhibition of the control. Mean values with different letters within a column are significantly different ( $P < 0.05$ ; LSD).

mainly attributed to its polyphenol content. These findings are in agreement with other reports showing that the antioxidant power and radical scavenging capacity of plant extracts were related to their high amount of polyphenols (56-61). Another study reported that the *Veronica officinalis* L. extract exhibited the most potent radical scavenging activity among other species. After characterization, it was found that the *V. officinalis* extract had a high amounts of phenolic when compared to other extracts (12). Moreover, in another experiment, anti-inflammatory and antioxidant activities of the *Veronica* spp. extract was supposed to be correlated with its polyphenols and iridoid compounds (62). Such direct relationship is a proof for the importance of these phytochemicals and their antioxidant activities.

Our *in vivo* results demonstrated the capability of the plant extract to improve the production of antioxidant enzymes in animal. SOD and CAT are antioxidant enzyme that provide a natural defense against oxidative stress of the cell (63). Treatment of animal models with carbon tetrachloride along with the plant extract showed an increase in the amounts of these enzymes in animal cells. Similarly, the extract increased the production of non-enzymatic antioxidants such as GSH. The thiobarbituric acid reactive substances (TBARS) assay was used for measuring the effects of the plant extract on lipid peroxidation. According to our results, groups treated with the plant extract showed a dose-dependent increase in the TBARS, thus indicating the ability of the extract in lipid peroxidation inhibition. These results are in agreement with recent experiments demonstrating the lipid peroxidation inhibitory potential of different plant extracts (64, 65). As known, the anti-lipid peroxidation activity by plant extracts can also be attributed to their polyphenol content. For example, the phenolic-rich extract of *Tetrapleura tetraptera* fruits showed effectively inhibited xanthine oxidase as well as  $F^{2+}$ -induced lipid peroxidation in animal models (66). Lipid peroxidation can generate a high amount of ROS, which, in turn, can be involved in cardiovascular diseases, diabetes mellitus, carcinogenesis, etc.(67). Therefore, the use of plants with high phenolic content and anti-lipid peroxidation activity might be beneficial for the prevention of many disorders.

Since oxidative stress is a deep-rooted cause of various disorders, the prevention of ROS and RNS production in the cell metabolism is of utmost importance. Vast usage of medicinal plants in different therapeutic contexts led us to study *V. persica* in order to evaluate its antioxidant and radical scavenging activities. Our results indicated the tremendous potential of this plant in reducing free radicals through different pathways, possibly due to its high polyphenol content. However, more investigations should be carried out to clarify the specific correlations between the plant bioactives and the observed biological activities.

### Acknowledgments

The authors are very grateful to Shahid Beheshti University of Medical Sciences, Tehran, Iran for financial support.

### Conflict of Interest

The authors declare no conflict of interest.

### References

1. Singh P, Mahadi F, Roy A, Sharma P. Reactive oxygen species, reactive nitrogen species and antioxidants in etiopathogenesis of diabetes mellitus type-2. *Indian Journal of Clinical Biochemistry* 2009; 24(4): 324-342.
2. Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *The International Journal of Biochemistry and Cell Biology* 2007; 39(1): 44-84.
3. Rice-evans CA, Miller NJ, Bolwell PG, Bramley PM, Pridham JB. The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radical Research* 1995; 22(4): 375-383.
4. Harput US, Saracoglu I, Inoue M, Ogihara Y. Phenylethanoid and iridoid glycosides from *Veronica persica*. *Chemical and Pharmaceutical Bulletin* 2002; 50(6): 869-871.
5. Sharifi-Rad J, Ayatollahi SA, Varoni EM *et al.* Chemical composition and functional properties of essential oils from *Nepeta schiraziana* Boiss. *Farmacia* 2017; 65(5): 802-812.
6. Zal F, Taheri R, Khademi F, Keshavarz E, Rajabi S, Mostafavi-Pour Z. The combined effect of furosemide and propranolol on GSH homeostasis in ACHN renal cells. *Toxicology Mechanisms and Methods* 2014; 24(6): 412-416.
7. Velioglu Y, Mazza G, Gao L, Oomah B. Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *Journal of Agricultural and Food Chemistry* 1998; 46(10): 4113-4117.
8. Amarowicz R, Pegg R, Rahimi-Moghaddam P, Barl B, Weil J. Free-radical scavenging capacity and antioxidant activity of selected plant species from the *Canadian prairies*. *Food Chemistry* 2004; 84(4): 551-562.
9. Sharifi-Rad J, Sureda A, Tenore GC *et al.* Biological activities of essential oils: From plant chemoeology to traditional healing systems. *Molecules* 2017; 22(1): 70.
10. Pratt D. Natural antioxidants from plant material. *ACS Publications*; 1992.
11. Sharifi Rad J, Hoseini Alfatemi SM, Sharifi Rad M, Iriti M. Free radical scavenging and antioxidant activities of different parts of *Nitraria schoberi* L. *Journal of Biologically Active Products from Nature* 2014; 4(1): 44-51.
12. Harput US, Genç Y, Khan N, Saracoglu I. Radical scavenging effects of different *Veronica* species. *Records of Natural Products* 2011; 5(2): 100.
13. Salehi B, Ayatollahi S, Segura-Carretero A *et al.* Bioactive chemical compounds in *Eremurus persicus* (Joub. & Spach) Boiss. essential oil and their health implications. *Cellular and Molecular Biology (Noisy-le-Grand, France)* 2017; 63(9): 1-7.
14. Saracoglu I, Harput US, Inoue M, Ogihara Y. New phenylethanoid glycosides from *Veronica pectinata* var. *glandulosa* and their free radical scavenging activities. *Chemical and Pharmaceutical Bulletin* 2002; 50(5): 665-668.
15. Harput US, Saracoglu I, Inoue M, Ogihara Y. Anti-inflammatory and cytotoxic activities of five *Veronica* species. *Biological and Pharmaceutical Bulletin* 2002; 25(4): 483-486.
16. Chari VM, Grayer-Barkmeijer RJ, Harborne JB, Österdahl B-G. An acylated allose-containing 8-hydroxyflavone glycoside from *Veronica filiformis*. *Phytochemistry* 1981; 20(8): 1977-1979.
17. Wolfe K, Wu X, Liu RH. Antioxidant activity of apple peels. *Journal of Agricultural and Food Chemistry* 2003; 51(3): 609-614.
18. Ordonez A, Gomez J, Vattuone M. Antioxidant activities of *Sechium edule* (Jacq.) Swartz extracts. *Food chemistry* 2006; 97(3): 452-458.
19. Kumaran A, Karunakaran RJ. In vitro antioxidant activities of methanol extracts of five *Phyllanthus* species from India. *LWT-Food*

Science and Technology 2007; 40(2): 344-352.

20. Sun J-S, Tsuang Y-H, Chen I-J, Huang W-C, Hang Y-S, Lu F-J. An ultra-weak chemiluminescence study on oxidative stress in rabbits following acute thermal injury. *Burns* 1998; 24(3): 225-231.

21. Liyana-Pathirana CM, Shahidi F. Antioxidant activity of commercial soft and hard wheat (*Triticum aestivum* L.) as affected by gastric pH conditions. *Journal of Agricultural and Food Chemistry* 2005; 53(7): 2433-2440.

22. Garrat D. *The Quantitative Analysis of Drugs*, Chapman and Hall Ltd. London; 1964.

23. Ruch RJ, Cheng S-j, Klaunig JE. Prevention of cytotoxicity and inhibition of intercellular communication by antioxidant catechins isolated from Chinese green tea. *Carcinogenesis* 1989; 10(6): 1003-1008.

24. Ng T, Liu F, Lu Y, Cheng C, Wang Z. Antioxidant activity of compounds from the medicinal herb *Aster tataricus*. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology* 2003; 136(2): 109-115.

25. Sharifi-Rad J, Miri A, Hoseini-Alfatemi SM, Sharifi-Rad M, Setzer WN, Hadjiakhoondi A. Chemical composition and biological activity of *Pulicaria vulgaris* essential oil from Iran. *Natural Product Communications* 2014; 9(11): 1633-1636.

26. Misra HP, Fridovich I. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *Journal of Biological Chemistry* 1972; 247(10): 3170-3175.

27. Niehaus W, Samuelsson B. Formation of malonaldehyde from phospholipid arachidonate during microsomal lipid peroxidation. *The FEBS Journal* 1968; 6(1): 126-130.

28. Pari L, Latha M. Protective role of *Scoparia dulcis* plant extract on brain antioxidant status and lipidperoxidation in STZ diabetic male Wistar rats. *BMC Complementary and Alternative Medicine* 2004; 4(1): 16.

29. Ellman GL. Tissue sulfhydryl groups. *Archives of Biochemistry and Biophysics* 1959; 82(1): 70-77.

30. Sharifi-Rad J, Salehi B, Stojanović-Radić ZZ *et al.* Medicinal plants used in the treatment of tuberculosis-Ethnobotanical and ethnopharmacological approaches. *Biotechnology Advances* 2017; doi: 10.1016/j.biotechadv.2017.07.001.

31. Sharifi-Rad J, Mnayer D, Tabanelli G *et al.* Plants of the genus *Allium* as antibacterial agents: From tradition to pharmacy. *Cellular and Molecular Biology (Noisy-le-Grand, France)* 2016; 62(9): 57-68.

32. Sharifi-Rad M, Tayeboon G, Miri A *et al.* Mutagenic, antimutagenic, antioxidant, anti-lipoxygenase and antimicrobial activities of *Scandix pecten-veneris* L. *Cellular and Molecular Biology (Noisy-le-Grand, France)* 2016; 62(6): 8-16.

33. Sharifi-Rad M, Iriti M, Gibbons S, Sharifi-Rad J. Anti-methicillin-resistant *Staphylococcus aureus* (MRSA) activity of Rubiaceae, Fabaceae and Poaceae plants: A search for new sources of useful alternative antibacterials against MRSA infections. *Cellular and Molecular Biology (Noisy-le-Grand, France)* 2016; 62(9): 39-45.

34. Sharifi-Rad J, Fallah F, Setzer W, Entezari RH, Sharifi-Rad M. *Tordylium persicum* Boiss. & Hausskn extract: A possible alternative for treatment of pediatric infectious diseases. *Cellular and Molecular Biology (Noisy-le-Grand, France)* 2016; 62(9): 20-26.

35. Salehi B, Mishra AP, Shukla I *et al.* Thymol, thyme and other plant sources: health and potential uses. *Phytotherapy Research* 2018 ; doi: 10.1002/ptr.6109.

36. Sharifi-Rad M, Mnayer D, Flaviana Bezerra Morais-Braga M *et al.* *Echinacea* plants as antioxidant and antibacterial agents: From traditional medicine to biotechnological applications. *Phytotherapy Research* 2018; doi: 10.1002/ptr.6101.

37. Sharifi-Rad M, Varoni EM, Iriti M *et al.* Carvacrol and Human Health: A Comprehensive Review. *Phytotherapy Research* 2018;

doi: 10.1002/ptr.6103.

38. Kristanc L, Kreft S. European medicinal and edible plants associated with subacute and chronic toxicity part I: Plants with carcinogenic, teratogenic and endocrine-disrupting effects. *Food and Chemical Toxicology* 2016; 92: 150-164.

39. Salehi B, Kumar NVA, Şener B, Sharifi-Rad M, Kılıç M, Mahady GB, Vlasisavljevic S *et al.* Medicinal Plants Used in the Treatment of Human Immunodeficiency Virus. *International Journal of Molecular Sciences* 2018; 19(5): 1459.

40. Cho EJ, Um SI, Han JH *et al.* The cytoprotective effect of *Rumex aquaticus* Herba extract against hydrogen peroxide-induced oxidative stress in AGS cells. *Archives of Pharmacal Research* 2016; 39(12): 1739-1747.

41. Sharifi-Rad M VE, Salehi B, Sharifi-Rad J, Matthews KR, Aya-tollahi SA, Kobarfard F, Ibrahim SA, Mnayer D, Zakaria ZA, Sharifi-Rad M, Yousaf Z, Iriti M, Basile A, Rigano D. . Plants of the Genus *Zingiber* as a Source of Bioactive Phytochemicals: From Tradition to Pharmacy. *Molecules* 2017; 22(12): 2145.

42. Salehi B, Zucca P, Sharifi-Rad M *et al.* Phytotherapeutics in cancer invasion and metastasis. *Phytotherapy Research* 2018; doi:10.1002/ptr.6087.

43. Sharifi-Rad J, Salehi B, Varoni EM *et al.* Plants of the *Melaleuca* genus as antimicrobial agents: from farm to pharmacy. *Phytotherapy Research* 2017; 31(10): 1475-1494.

44. Sharifi-Rad J, Salehi B, Schnitzler P *et al.* Susceptibility of herpes simplex virus type 1 to monoterpenes thymol, carvacrol, p-cymene and essential oils of *Sinapis arvensis* L., *Lallemantia royleana* Benth. and *Pulicaria vulgaris* Gaertn. *Cellular and Molecular Biology (Noisy le Grand)* 2017; 63(8): 42-47.

45. Sharifi-Rad J, Mnayer D, Roointan A *et al.* Antibacterial activities of essential oils from Iranian medicinal plants on extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli*. *Cellular and Molecular Biology (Noisy-le-Grand, France)* 2016; 62(9): 75-82.

46. Stojanović-Radić Z, Pejić M, Stojanović N, Sharifi-Rad J, Stanković N. Potential of *Ocimum basilicum* L. and *Salvia officinalis* L. essential oils against biofilms of *P. aeruginosa* clinical isolates. *Cellular and Molecular Biology (Noisy-le-Grand, France)* 2016; 62(9): 27-32.

47. Sharifi-Rad M, Tayeboon G, Sharifi-Rad J, Iriti M, Varoni E, Razazi S. Inhibitory activity on type 2 diabetes and hypertension key-enzymes, and antioxidant capacity of *Veronica persica* phenolic-rich extracts. *Cellular and Molecular Biology (Noisy-le-Grand, France)* 2016; 62(6): 80-85.

48. Bagheri G, Mirzaei M, Mehrabi R, Sharifi-Rad J. Cytotoxic and Antioxidant Activities of *Alstonia scholaris*, *Alstonia venenata* and *Moringa oleifera* Plants From India. *Jundishapur Journal of Natural Pharmaceutical Products* 2016; 11(3): e31129.

49. Sahraie-Rad M, Izadyari A, Rakizadeh S, Sharifi-Rad J. Preparation of strong antidandruff shampoo using medicinal plant extracts: a clinical trial and chronic dandruff treatment. *Jundishapur Journal of Natural Pharmaceutical Products* 2015; 10(4): e21517.

50. Bojor O. *Guide of medicinal and aromatic plants from A to Z. Romanian: Ghidul plantelor medicinale și aromatice de la A la Z: Fiat Lux Bucharest*; 2003.

51. Fierascu RC, Georgiev MI, Fierascu I *et al.* Mitodepressive, antioxidant, antifungal and anti-inflammatory effects of wild-growing Romanian native *Arctium lappa* L.(Asteraceae) and *Veronica persica* Poiret (Plantaginaceae). *Food and Chemical Toxicology* 2017.

52. Vogl S, Picker P, Mihaly-Bison J *et al.* Ethnopharmacological in vitro studies on Austria's folk medicine - An unexplored lore in vitro anti-inflammatory activities of 71 Austrian traditional herbal drugs. *Journal of Ethnopharmacology* 2013; 149(3): 750-771.

53. Mocan A, Vodnar DC, Vlase L, Crişan O, Gheldiu A-M, Crişan G. Phytochemical characterization of *Veronica officinalis* L., V. teu-

- cium L. and *V. orchidea* Crantz from Romania and their antioxidant and antimicrobial properties. *International Journal of Molecular Sciences* 2015; 16(9): 21109-21127.
54. Stojković DS, Živković J, Soković M et al. Antibacterial activity of *Veronica montana* L. extract and of protocatechuic acid incorporated in a food system. *Food and Chemical Toxicology* 2013; 55: 209-213.
55. Živković J, Barreira JC, Stojković D et al. Phenolic profile, antibacterial, antimutagenic and antitumour evaluation of *Veronica urticifolia* Jacq. *Journal of Functional Foods* 2014; 9: 192-201.
56. Hanganu D, Filip L, Olah N-K et al. Evaluation of polyphenolic profile and antioxidant activity for *Cytisus nigricans* and *Cytisus albus*. *Farmacia* 2016; 64(6): 863-867.
57. de Beer D, Miller N, Joubert E. Production of dihydrochalcone-rich green rooibos (*Aspalathus linearis*) extract taking into account seasonal and batch-to-batch variation in phenolic composition of plant material. *South African Journal of Botany* 2017; 110: 138-143.
58. Perera D, Soysa P, Wijeratne S. Polyphenols contribute to the antioxidant and antiproliferative activity of *Phyllanthus debilis* plant in-vitro. *BMC Complementary and Alternative Medicine* 2016; 16(1): 339.
59. Peng Y, Zhang H, Liu R et al. Antioxidant and anti-inflammatory activities of pyranoanthocyanins and other polyphenols from staghorn sumac (*Rhus hirta* L.) in Caco-2 cell models. *Journal of Functional Foods* 2016; 20: 139-147.
60. Nair AC, Goveas S, D'Souza L, D'Cunha F, D'Souza V. Antibacterial and antioxidant potential of organic solvents extract of mangrove endophytic fungus *Eupenicillium senticosum* Scott. *Journal of Alternative Medicine Research* 2017; 9(1): 65.
61. Hashemi SMB, Brewer MS, Safari J et al. Antioxidant activity, reaction mechanisms, and kinetics of *Matricaria recutita* extract in commercial blended oil oxidation. *International Journal of Food Properties* 2016; 19(2): 257-271.
62. Dussossoy E, Brat P, Bony E et al. Characterization, anti-oxidative and anti-inflammatory effects of Costa Rican Noni Juice (*Morinda citrifolia* L.). *Journal of Ethnopharmacology* 2011; 133(1): 108-115.
63. Abed HN, Hadwan MH. Clinical and thermodynamical properties of catalase enzyme in sera of Iraqi diabetic patients. *BioTechnology: An Indian Journal* 2016; 12(1).
64. Irondi EA, Oboh G, Agboola SO, Boligon AA, Athayde ML. Phenolics extract of *Tetrapleura tetraptera* fruit inhibits xanthine oxidase and Fe<sup>2+</sup>-induced lipid peroxidation in the kidney, liver, and lungs tissues of rats *in vitro*. *Food Science and Human Wellness* 2016; 5(1): 17-23.
65. Thayyil AH, Muthu AK, Ibrahim M. In vivo antioxidant and lipid peroxidation effect of various extracts from aerial parts of *Chomelia asiatica* (Linn) in rat fed with high fat diet. *African Journal of Pharmacy and Pharmacology* 2016; 10(38): 810-816.
66. Irondi EA, Oboh G, Agboola SO, Boligon AA, Athayde ML. Phenolics extract of *Tetrapleura tetraptera* fruit inhibits xanthine oxidase and Fe<sup>2+</sup>-induced lipid peroxidation in the kidney, liver, and lungs tissues of rats *in vitro*. *Food Science and Human Wellness* 2016; 5(1): 17-23.
67. Ngoc TM, Khoi NM, Nhiem NX et al. Xanthine oxidase inhibitory activity of constituents of *Cinnamomum cassia* twigs. *Bioorganic and Medicinal Chemistry Letters* 2012; 22(14): 4625-4628.