



Original Research

## Importance of miR-125a-5p and miR-122-5p expression in patients with HBV infection

Ramin Lak<sup>1,2</sup>, Ramin Yaghobi<sup>3\*</sup>, Masoud Garshasbi<sup>4</sup>

<sup>1</sup> Department of Microbiology, Fars Science and Research Branch, Islamic Azad University, Fars, Iran

<sup>2</sup> Department of Microbiology, Shiraz branch, Islamic Azad University, Shiraz, Iran

<sup>3</sup> Shiraz Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>4</sup> Department of Medical Genetics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

\*Correspondence to: [rayaviro@yahoo.com](mailto:rayaviro@yahoo.com)

Received October 22, 2019; Accepted June 22, 2020; Published July 31, 2020

Doi: <http://dx.doi.org/10.14715/cmb/2020.66.5.1>

Copyright: © 2020 by the C.M.B. Association. All rights reserved.

**Abstract:** MicroRNAs (miRNAs) as a small RNA and post-transcriptional modulator are shown to have regulatory effects for different cellular activities and pathways, such as metabolism, virus replication and also cell growth. In addition, miRNAs can regulate the replication of the hepatitis B virus (HBV). Therefore, the expression profile of miRNAs was evaluated in HBV infected patient groups and healthy controls. The expression levels of following microRNAs (as noninvasive biomarkers) were compared in two experimental (those with various stages of HBV infection) and control groups to evaluate their diagnosis ability: miR-122-5p, miR-125a-5p, miR-199a-3p, miR-210-5p, miR-205-5p, miR-155-5p, miR-372-5p, and miR-1-5p. RNA extraction was performed for 45 serum samples. The miRCURY LNA™ Universal RT-miRNA-PCR system and miRNA PCR panels were used for measuring microRNA expression profiles. To normalize quantitative values, the endogenous reference by UniSp6 expression was used. Serum miR-125a-5p and miR-122-5p were significantly higher in patients in different stages of HBV infection ( $p < 0.001$ ) than in controls ( $p < 0.001$ ). Receiver operating characteristic (ROC) curve analyses suggested that serum miR-125a-5p and miR-122-5p have significant diagnostic value for HBV infection. A significant difference was not found in terms of serum levels of other miRs (miR-199a-3p, miR-210-5p, miR-205-5p, miR-155-5p, miR-372-5p and miR-1-5p). Results suggest that miR-125a-5p and miR-122-5p can be used as possible noninvasive biomarkers for monitoring of HBV infection need to confirm in future completed studies.

**Key words:** MicroRNAs; Hepatitis B Virus; Biomarker.

### Introduction

Studies have shown that one-third of the world's population is influenced by Hepatitis B virus (HBV) infection, which is considered a certain health concern all over the world, of whom over 350 million people have diagnosed with chronic HBV (CHB) infection (1). HBV infection can be associated with several complications, such as asymptomatic carriers, liver cirrhosis and hepatocellular cancer (2,3). Therefore, understanding the mechanism underlying chronic HBV infection at the molecular level is of great importance for the optimal management of chronic HBV infection. MicroRNAs (miRNAs) have shown to possibly be associated with HBV infection regulation and relevant disorders (3, 4).

It has indicated that miRNAs as small endogenous noncoding RNAs containing 21–25 nucleotides are capable of post-transcriptional gene expression regulation via target mRNAs degradation induction or inhibition of their translation into protein (5). Recent several studies indicated that miRNAs might affect virus-host interactions and play a crucial role in the viral life cycle and pathogenesis (6–8). Indeed, HBV replication can be regulated by miRNAs directly through linking to the HBV genome or indirectly via modulating transcriptional factors linked to HBV for the regulation of HBV enhan-

cer/promoter activities (9, 10). Zhang et al. were first studied miRNAs, which directly target HBV transcripts. They found that HBsAg expression and HBV replication were appropriately decreased by miR-199a-3p and miR-210-5p (11). In another study, the authors stated that HBV replication is promoted by miRNA-1 via an elevation in transcriptional activity of HBV core promoter (12). MiRNAs not only are interacted with HBV transcripts, they have also regulated immune responses and signaling pathways, so they are essential for HBV replication and pathogenesis (13, 14). MiR-155-5p has shown to increase the expression level of Interferon-inducible antiviral genes against HBV via direct prevention of suppressor of cytokine signaling 1 (SOCS1) expression levels, which results in an elevation in JAK/STAT signaling pathway (14).

Recognizing the appropriate non-invasive biomarkers for liver disease caused by the HBV virus is challenging, which can lead to the early effective treatment and also desirable treatment outcomes (15). Novel methods have emerged to use miRNAs as biomarkers for liver disease by recognition of miRNAs capable of indicating pathophysiological conditions of human bodily fluids (HBF). In addition, they are considered as suitable biomarkers for early and pre-symptomatic diagnosis due to their availability, great stability in the circulation and also expression patterns consistent with the tissues (16).

Serum miRNA profiles from HBV carriers for recognizing appropriate biomarkers have widely investigated (3,13–17). Therefore, in this study, the expression profile of a panel of miRNAs was analyzed in sera of HBV infected patient groups and healthy controls.

## Materials and Methods

### Study population

The present research was performed at Pars Hospital Laboratory, in Tehran, Iran from January 2015 to January 2017. All admitted untreated patients with CHB (n=45) based on immunological results divided into three groups: HBeAg positive, HBeAg negative, OHB and healthy controls (n=15) were recruited. CHB diagnosed by serum HBs antigen for more than 6 months was considered as the inclusion criterion. Patients exhibiting the following were excluded from the study: alcoholic liver disease, organ transplantation, immunosuppression and malignant comorbidities within the last 5 years. All participants gave written informed consent.

### Serum sample collection

Blood samples were collected from each participant and kept on ice followed by centrifugation at 1500× g for 10 min at 4°C. The resulting supernatant was sent to three RNase/DNase-free tubes separately (Immunological-biochemical, HBV Molecular testing and miRNA study) and was stored at -80°C until analysis.

### Clinical Chemistry and immunological markers

Cobas Integra 400 Plus (Roche Products Ltd., Switzerland) was used for serum ALT and AST levels according to the manufacturer's instructions. The levels of HBsAg, HBeAg, anti-HBeAg, anti-HBs, anti-HBc, and AFP were tested by automated available immunoassays (Roche Cobas e 411 analyzer).

### DNA isolation and RT-q-PCR for HBV

Magcore Viral Nucleic Acid Extraction Kit was used for HBV DNA extraction from 400 µl of each serum sample. Flurion HBV QNP 2.0 PCR kit (Iontek, Turkey) was also employed for real-time PCR on Bio-Rad CFX96. The following steps were considered for amplification: Initial hot-start denaturation (95°C/15 min), followed by 50 cycles (95°C/30 s), annealing and extension (54°C/90 s). Genotypes of HBV were determined using AmpliSens® HBV genotype-FRT PCR kit variant FRT by real-time PCR. The following steps were considered for amplification: Initial hot-start denaturation (95°C/2 min), followed by 40 cycles of denaturation (95°C/25 s), annealing (58°C/25 s), extension (72°C/1 min), and final extension (72°C/5 min).

### miRNA analysis

#### RNA extraction and cDNA synthesis

200 µL of the serum was used to extract miRNA using a miRCURY™ RNA Isolation Kit-Biofluid (Exiqon, USA) based on the manufacturer's protocol. The optical density in 260/280nm was measured for assessing the purity and concentration of RNA. Agarose gel electrophoresis (1%) was applied in order to measure RNA integrity. cDNA synthesis was performed by miRCURY LNA™ Universal RT-cDNA-Synthesis

(Exiqon, USA), based on the provided protocol by the manufacturer.

### SYBR-Green Real-Time PCR

The level of gene expression was determined via CFX96 Real-Time thermal cycler (Biorad, USA). Real-time PCR for the 8 miRNAs (miR-122-5p, miR-125a-5p, miR-199a-3p, miR-210-5p, miR-205-5p, miR-155-5p, miR-372-5p, and miR-1-5p) were performed using miRNA PCR panels, miRCURY LNA™ Universal RT-miRNA-PCR system. The specific microRNA LNA™ PCR primers (Exiqon, Denmark) were used. To normalize the results, we were used UniSp6 as an internal standard control and all samples were run in duplicate. The reaction mixture was prepared by SYBR Green Premix (5µl), water (1µl) and 4µl of the control cDNA. The amplification of the reaction mixture was performed as follows: one cycle (95°C/10 min) followed by 40 cycles (95°C/10 s, 60°C/60 s, and 95°C/60 s, respectively). Melting Curve Analysis was used to prove the specificity of the amplified product.

### Statistics

Results are represented as the mean ± standard deviation (SD). One-way ANOVA and the Kruskal-Wallis test were used for comparisons between groups. Data were analyzed by SPSS 23 (IBM Corporation, USA). The p values of less than 0.05 were considered significant. For estimation of diagnostic ability of miRNA receiver operating characteristic (ROC) curve analysis was used via medcalc statistical software (Belgium).

## Results

### Clinical details of patients

The study included 45 patients with Hepatitis B infection and 15 healthy control. Patients with HBV infection were divided into 3 groups based on immunological markers. 15 patients were HBeAg positive chronic hepatitis, 15 patients were HBeAg negative chronic hepatitis and 15 patients were occult hepatitis B. Mean age of patients group was 38.04±8.9 SD (range 22-56 years) and the distribution of gender in the patient group was 36 males (73.3%) and 9 female (%26.6). The mean age of the control group was 33.74±5.7 SD (range 26-44 years) and the distribution of gender in the control group was 9 male (%60) and 6 females (%40).

Separately in different patient groups, the mean age of ENCH Patient was 36.42±7.9 SD (range 27-49 years) and the distribution of gender in ENCH group was 12 males (80%) and 3 females (%20). The mean age of EPCH Patient was 36.63±8.3 SD (range 22-47 years) and the distribution of gender in the EPCH group was 11 males (73.3%) and 4 females (%26.6). The mean age of OHB Patient was 41.5±7.8 SD (range 29-56 years) and the distribution of gender in the OHB group was 12 male (80%) and 3 females (%20). The crowding index was successfully determined in patients and healthy groups. In index value <0.5; just 13.3% of control groups. In index value 0.5-0.99; 1.5% were control group, 6.6% were EPCH group, 13.3% were ENCH group and 6.6% were OHB. In index value >1; 73.3% were control group, 93.3% were EPCH and 93.3% were OHB.

## Biochemical and molecular markers and HBV groups

All patients were negative for anti-HCV, anti-HIV+Ag and anti-Hbc-IgM. Mean ALT, AST and AFP of patients was  $29.24 \pm 18.3$  SD,  $26.18 \pm 15.3$  SD and  $7.17 \pm 15.0$  SD respectively. Whereas mean ALT, AST and AFP OF Controls was  $25.80 \pm 7.94$  SD,  $23.70 \pm 5.78$  SD and  $2.12 \pm 0.56$  SD respectively. There was no statistically significant difference between the patients and the control group in this value. But mean ALT, AST and AFP of ENCH group was:  $27.2 \pm 14.46$  SD,  $24.60 \pm 11.4$  SD and  $5.97 \pm 10.4$  SD, EPCH group:  $45.6 \pm 25.5$  SD,  $37.53 \pm 21.71$  SD and  $6.35 \pm 11.3$  SD, OHB group:  $20.93 \pm 3.12$  SD,  $19.40 \pm 2.4$  SD and  $2.4 \pm 0.69$ . The means of ALT and AST were significantly higher ( $p < 0.001$ ) in the EPCH group than in ENCH and OHB and healthy controls. The positive cases of HBsAb in control, EPCH, ENCH and OHB were 66.7%, 0%, 0% and 53.3% respectively. The positive cases of HBsAb in control, EPCH, ENCH and OHB were 5%, 100%, 100% and 100% respectively. The positive cases of HBsAg were 0%, 100%, 0% and 0%, in HBeAb were 0%, 0%, 100% and 100% and the positive cases of HBsAg were 0%, 100%, 100% and 0% respectively. All patients were positive for HBV Nested PCR and Taqman real-time PCR Methods simultaneously. HBV genotypes were successfully determined in all of the patients and 100% were genotype D. The EPCH cases had high HBV viral load ( $3.5 \pm 0.7$ ) than in ENCH patients (Mean  $7.1E+05$ ) and OHB cases (mean  $8.3.E+02$ ). The results of many clinical tests are quantitative and are provided on a continuous scale.

To help decide the presence or absence of disease, a


cut-off point is chosen and results that are below the cut-off are regarded as normal. The receiver operating characteristic (ROC) curve is widely accepted as a method for selecting an optimal cut-off point and for comparing the accuracy of diagnostic testing. The ROC curve is also important because the area under the curve (AUC) is a reflecting of how good the test is at distinguishing between the patient with the disease and without disease. A perfect test will have an AUC of 1.0, while a completely useless test has AUC of 0.5. The AUC of many clinical tests falls between these two values one way of interpreting the AUC is that a test with an area greater the 0.9 has high accuracy, while 0.7-0.9 indicates moderate accuracy, 0.5-0.7 low accuracy and 0.5 a chance result. In order to check the accuracy of the tests done in the thesis and their ability to detect hepatitis B infection, we used the ROC curve analysis. The result of different performed diagnostic tests is shown in Fig. 1 & 2.

## MiRNA profile and HBV groups




The serum levels of miRNA including: miR-122-5p, miR-125a-5p, miR-199a-3p, miR-210-5p, miR-205-5p, miR-155-5p, miR-372-5p, and miR-1-5p) were analyzed in different samples from ENCH, EPCH, OHB and healthy control groups.

The serum miRNA levels in different Patient group including: ENCH, EPCH and OHB was not correlated significantly with each other. The serum level of miR-122-5p, miR-125-5p, and residual miRNAs were significantly different between all patient groups and healthy control ( $p < 0.001$ ) (Fig. 3 & 4).

The heat map analysis is presented in Fig.5. Data




**Figure 1.** ROC curve analysis of immunological and molecular testing of HBV (Medcalc software).



with increase/decrease of expression are highlighted red/blue, respectively and unchanged data are displayed white. As shown on the side of the heat map, color intensity was calibrated to the expression level. 2 miRNAs from 8 assessed miRNAs showed differential expression in HBV infected patients compared to the control groups. The up-regulated miRNAs included miR-122-5p and miR-125-5p, which implies the important role of their binding sites in aberrant miRNA expression caused by HBV in the host target genes.

**Discussion**

The role of miRNAs in HBV infection has attracted great interest in recent years, since accumulating evidence has suggested that miRNAs play crucial roles in the HBV pathogenesis (3, 4, 9). Cirrhosis and hepatocellular carcinoma (HCC) known for their high mortality rate are among the complications of HBV infection (18,19). The present study aimed at assessing the serum levels of eight miRs, including miR-122-5p, miR-125a-5p, miR-199a-3p, miR-210, miR-205, miR-155, miR-372 and miR-1 in patients infected with HBV in clini-



**Figure 5.** Heatmap of miR levels in different cases related to their group.

cal different stages compared with the healthy control group.

The present study, which supported the results reported by previous studies, showed up-regulated expression of miR-125a-5p in HBV patients compared with the healthy control groups. The prediction of target sites for human microRNAs within the HBV genome was first made by Potenza et al. and they indicated that miR-125a-5p can bind to the target sequences in the viral genome. Indeed, miR-125-5p can affect HBsAg expression, as miR-125a mimic or inhibitor transfection leading to secretion of HBsAg by PLC/PRF/5 cells induced a noticeable reduction or increase in HBsAg expression, respectively (20). In the HBV-producing cell line HepG2.2.15, hsa-miR-125a-5p expression was indicated at least three times more than parental HepG2 cells (21). Moreover, authors show that treatment with iron can lead to a reduced level of hsa-miR-125a-5p expression via an enhancement in HBV replication, whereas by TGF- $\beta$  treatment, an elevation can be seen in the expression level of hsa-miR-125a-5p due to the decreased HBV replication (22). Accordingly, HBV replication can be prevented by miR-125a-5p via direct binding to viral transcripts and the HBV sequence targeted by hsa-miR125a-5p encodes amino acids falling within an important segment (for cell binding) of the extracellular pre-S1 domain of HBsAg (23).

Recently, miR-122, miR-130, miR-183, miR-196, miR-209 and miR-96 were presented as potential biomarkers to distinguish different stages of hepatitis and liver injuries (24). However, there are conflicting results regarding the relationship between miRNA-122-5p expression levels and HBV replication. In this study, we have found a significant difference in the serum level of miRNA-122-5p among all patient groups and healthy control ( $p < 0.001$ ). Consistent with our results, Ebrahimi-fard et al indicated that the rate of miRNA-122-5p expression in patients with CHB and HBV related cirrhosis was 1.8 times significantly more than the control group ( $P < 0.05$ ) (25). In contrast, Jian Zhou et al. indicated

that miRNA-122-5p expression levels in HBV patients were decreased (26). Bandopadhyay et al. compared the expression level of miRNA-122-5p in HepG2.2.15 with HBV-expressing cell line and HepG2 unstable situation. They found down-regulation of miRNA-122-5p in HepG2.2.15 compared to controls HepG2 (27). In a rat model miR-511-3p in plasma was recommended as a possible biomarker for liver sinusoidal endothelial cells damage (28). MiR-122-5p as the most abundant microRNA in the liver plays an important role in biological functions and significance in various diseases (29). Although the exact role of miRNA-122-5p in HBV infection is unclear, miRNA-122-5p can bind directly to the polymerase region of viral transcripts, which can result in inhibition of HBV gene expression and replication. HBV replication can also be suppressed by MiRNA-122-5p via Cyclin G1 down-regulation as well as an increase in p53 activities. It can be concluded that hepatocyte-specific miRNA-122-5p is a vital regulator of HBV replication via direct expression of viral RNA or by modulating the expression of the host gene (30).

In this study, we found no significant difference between the six miRs (miR-199a-3p, miR-210, miR-205, miR-155, miR-372, and miR-1) in the serum of patients infected with HBV compared to the control group. Conflicting results might be due to the difference in the participants' age, samples (peripheral blood mononuclear cells (PBMCs), liver biopsy or serum) and the environmental factors (e.g., alcoholic liver disease, etc.). Our data indicated that mean ALT and AST were significantly higher ( $p < 0.001$ ) in the EPCH group than in ENCH and OHB and healthy controls, suggesting a positive correlation with hepatic necroinflammation.

ROC curve analysis was employed to assess whether serum miRs can be used as a diagnostic marker for CHB and OHB. Its findings indicated that miRNA-122-5p and miR-125a-5p serum levels were useful markers for discriminating patients infected HBV. An AUC of ROC curve for miR-125a-5p is 0.778 (Fig. 5d), with the sensitivity and specificity 77.8 and 100.0 %, respectively. For

miR-122-5p, an AUC of ROC curve is 0.822 (Fig. 5b), with a sensitivity/specificity ratio of 82.2/100.0 %. This suggests that miRNAs can clearly improve the diagnostic accuracy of efficacy evaluation in HBV infection. In addition to, The AUC for HBcAb (1.00,  $P < 0.001$ ) was equal to HBV viral load (1.00,  $P < 0.001$ ) and markedly higher than that of HBsAg (0.86,  $P < 0.001$ ), HBeAb (0.83,  $P < 0.001$ ), ALT (0.74,  $P < 0.001$ ), HBeAg (0.66,  $P < 0.001$ ), and HBsAb (0.6,  $P = 0.001$ ).

Cellular miRNAs can influence HBV replication via following ways; 1) they can bind to HBV transcripts, 2) via affecting cellular factors related to the life cycle of HBV, 3) through regulating genes and signaling pathways essential for pathogenesis of HBV, 4) via adjusting epigenetic reformations, including histone modification and methylation (15,31). MiRNAs can directly interact with viral components, which has empirically been approved, including miR199a-3p (affecting HBsAg coding region) (11), miR210 (affecting HBV pre-S1 region) (11) and miR205 (affecting X gene) (32). Replication of HBV is effectively suppressed by MiR-141 and miR-155 directly by affecting peroxisome proliferator-activated receptor alpha (PPAR $\alpha$ ) as well as CAAT Enhancer binding protein  $\beta$  (C/EBP- $\beta$ ), respectively (33). However, it has indicated that the replication of HBV is promoted by miR-1, miR-372, and miR-501 by influencing host gene expression. MiR-1 as an example is able to alter the expression level of several genes via affecting histone deacetylase 4 (HDAC4), including up-regulation of farnesoid X receptor (FXR) leading to an increase in transcription and replication of HBV via binding to the HBV core promoter (12). MiR-372 is capable of increasing the expression level of HBV via NFIB transcription factor. NFIB levels were lower in a stable HBV-producing cell line (HepG2.2.15 cells) with elevated endogenous expression of miRs-372 in comparison with the control cell line (HepG2 cells) (34). There are limitations to research on gene expression that researchers need to consider (35-47). In these regards immunogenicity of the multi-epitopic recombinant glycoproteins virus: implications may be concerned (48). There are also several biotechnological research on the virus (49-61), each of which can be helpful in applying the results.

There are several limitations to this study. First, a larger sample size should be considered to prove the diagnostic characteristic of microRNAs. Second, nothing patients accepted liver biopsies; we did not measure the miRs expression in liver.

In conclusion, the increase in miR-122-5p and miRNA-125b-5p levels in serum of HBV infected patients compared with healthy normal controls were found. However; other miRs have not shown a significant difference among patients and controls. These results highlighted that further studies are needed on the effect of miRs as noninvasive serum biomarkers to diagnose HBV outcomes.

### Acknowledgments

This study was supported by Islamic Azad University. We appreciate all patients who participated in this study and thanks also from Dr. Afsson Afshari for her scientific and statistic advice. Thanks to Bistoon Shafa Co., Iran (website: www.camelina.info, Instagram: @came-

lina.ir) for all supports.

### Interest conflict

The authors have no conflict of interest to declare.

### Author contributions

Study concept and design: Lak R, Yaghobi R, Garshasbi M, Performed the experiments: Lak R, Drafting of the manuscript: Lak R, Yaghobi R, Garshasbi M. Statistical analysis: Lak R, Yaghobi R, Garshasbi M.

### References

- MacLachlan JH, Cowie BC. Hepatitis B virus epidemiology. *Cold Spring Harb Perspect Med* 2015;5.
- Liang TJ. Hepatitis B: The virus and disease. *Hepatology* 2009;49: S13-21.
- Farokhzadeh Z, Dehbidi S, Geramizadeh B, Yaghobi R, Malekhosseini SA, Behmanesh M, et al. Association of MicroRNA Polymorphisms With Hepatocellular Carcinoma in an Iranian Population. *Ann Lab Med* 2019;39:58.
- Wang G, Dong F, Xu Z, Sharma S, Hu X, Chen D, et al. MicroRNA profile in HBV-induced infection and hepatocellular carcinoma. *BMC Cancer* 2017;17:805.
- Khoshtinat Nikkhai S, Dorostkar R, Ranjbar S, Heydarzadeh H, Tat M, Ghalavand M, Farasat A, Hashemzadeh MS. Synergistic Effect of Expressed miR-128 and Puma Protein on Targeted Induction of Tumor Cell Apoptosis. *Iran J Biotechnol.* 2016; 14(3): 185-191.
- Skalsky RL, Cullen BR. Viruses, microRNAs, and Host Interactions. *Annu Rev Microbiol.* 2010;64:123-41.
- Naqvi AR, Shango J, Seal A, Shukla D, Nares S. Viral miRNAs alter host cell miRNA profiles and modulate innate immune responses. *Front Immunol* 2018;9:433.
- Bernier A, Sagan SM. The diverse roles of microRNAs at the host-virus interface. *Viruses.* 2018; 10:440.
- Onorato L, Russo A, Sagnelli C, Sagnelli E, Potenza N, Coppola N. Micro-RNAs in hepatitis B virus-related chronic liver diseases and hepatocellular carcinoma. *World J Hepatol* 2018;10:558-70.
- Wang Z, Wu Z, Huang P. The function of miRNAs in hepatocarcinogenesis induced by hepatitis B virus X protein. *Oncol Rep* 2017;38:652-64.
- Zhang G ling, Li Y Xuan, Zheng S qi, Liu M, Li X, Tang H. Suppression of hepatitis B virus replication by microRNA-199a-3p and microRNA-210. *Antiviral Res* 2010;88:169-75.
- Zhang X, Zhang E, Ma Z, Pei R, Jiang M, Schlaak JF, et al. Modulation of hepatitis B virus replication and hepatocyte differentiation by MicroRNA-1. *Hepatology* 2011;53:1476-1485.
- Jiang X, Kanda T, Wu S, Nakamura M, Miyamura T, Nakamoto S, et al. Regulation of microRNA by hepatitis B virus infection and their possible association with control of innate immunity. *World J Gastroenterol* 2014;20:7197-06.
- Zhu H, Geng Y, He Q, Li M. miRNAs regulate immune response and signaling during hepatitis C virus infection. *Eur J Med Res* 2018;23:19.
- Kitab B, Alj Hs, Ezzikouri S, Benjellou S. MicroRNAs as Important Players in Host-hepatitis B Virus Interactions. *J Clin Transl Hepatol* 2015;3:141-61.
- Fu Y, Wei X, Tang C, Li J, Liu R, Shen A, et al. Circulating microRNA-101 as a potential biomarker for hepatitis B virus-related hepatocellular carcinoma. *Oncol Lett* 2013;6:1811-15.
- Su C, Hou Z, Zhang C, Tian Z, Zhang J. Ectopic expression of microRNA-155 enhances innate antiviral immunity against HBV infection in human hepatoma cells. *Virol J* 2011;8:354.
- Zamor P, Delemos A, Gastrointestinal MR-J of, 2017 U. *Viral*

hepatitis and hepatocellular carcinoma: Etiology and Management. *J Gastrointest Oncol* 2017;8:229–42.

19. Behzad-Behbahani A, Mojiri A, Tabei SZ, Farhadi-Andarabi A, Pouransari R, Yaghobi R, et al. Outcome of hepatitis B and C virus infection on graft function after renal transplantation. *Transplant Proc* 2005;37:3045–47.

20. Potenza N, Papa U, Mosca N, Zerbini F, Nobile V, Russo A. Human microRNA hsa-miR-125a-5p interferes with expression of hepatitis B virus surface antigen. *Nucleic Acids Res* 2011;39:5157–63.

21. Liu Y, Wang C-M, Han P, Wang L, Cheng Y-Q, Xu D-P, et al. Altered expression profiles of microRNAs in a stable hepatitis B virus-expressing cell line. *Chin Med J (Engl)* 2009;122:10–4.

22. Park SO, Kumar M, Gupta S. TGF- $\beta$  and Iron Differently Alter HBV Replication in Human Hepatocytes through TGF- $\beta$ /BMP Signaling and Cellular MicroRNA Expression. Bouchard M, editor. *PLoS One* 2012;7:e39276.

23. Le Duff Y, Blanchet M, Sureau C. The Pre-S1 and Antigenic Loop Infectivity Determinants of the Hepatitis B Virus Envelope Proteins Are Functionally Independent. *J Virol* 2009;83:12443–51.

24. Musaddaq G, Shahzad N, Ashraf MA, Arshad MI. Circulating liver-specific microRNAs as noninvasive diagnostic biomarkers of hepatic diseases in human. *Biomarkers* 2019;24:103–9.

25. Ebrahimifard M, Zandi M, Moradi A. Evaluation of miR-122 level in chronic HBV infection and liver cirrhosis patients. *Arch Med Lab Sci* 2016;2:108-10.

26. Zhou J, Yu L, Gao X, Hu J, Wang J, Dai Z, et al. Plasma microRNA panel to diagnose hepatitis B virus-related hepatocellular carcinoma. *J Clin Oncol* 2011;29:4781–8.

27. Bandothyay M, Sarkar N, Datta S, Das D, Pal A, Panigrahi R, et al. Hepatitis B virus X protein mediated suppression of miRNA-122 expression enhances hepatoblastoma cell proliferation through cyclin G1-p53 axis. *Infect Agent Cancer* 2016;11:40.

28. Oda S, Takeuchi M, Akai S, Shirai Y, Tsuneyama K, Yokoi T. miRNA in Rat Liver Sinusoidal Endothelial Cells and Hepatocytes and Application to Circulating Biomarkers that Discern Pathogenesis of Liver Injuries. *Am J Pathol* 2018;188:916–28.

29. Thirion M, Ochiya T. Roles of microRNAs in the hepatitis B virus infection and related diseases. *Viruses* 2013;5:2690–703.

30. Mizuguchi Y, Takizawa T, Uchida E. Host cellular microRNA involvement in the control of hepatitis B virus gene expression and replication. *World J Hepatol* 2015;7:696–02.

31. Ghanbari R, Ravanshad M, Hosseini SY, Yaghobi R, Shahzamani K. Genotyping and infection rate of GBV-C among Iranian HCV-infected patients. *Hepat Mon* 2010;10:80–7.

32. Zhang T, Zhang J, Cui M, Liu F, You X, Du Y, et al. Hepatitis B Virus X Protein Inhibits Tumor Suppressor miR-205 through Inducing Hypermethylation of miR-205 Promoter to Enhance Carcinogenesis. *Neoplasia* 2015;15:1282-91.

33. Sarkar N, sciences RC-I journal of molecular, undefined. Hepatitis B virus infection, microRNAs and liver disease. *Int J Mol Sci* 2015;16:17746–62.

34. Guo H, Liu H, Mitchelson K, Rao H, Luo M, Xie L, et al. MicroRNAs-372/373 promote the expression of hepatitis B virus through the targeting of nuclear factor I/B. *Hepatology* 2011;54:808–19.

35. Bordbar M, Darvishzadeh R, Pazhouhandeh M, Kahrizi D. An overview of genome editing methods based on endonucleases. *Modern Genetics J* 2020; 15(2): 75-92.

36. Ghaheri M, Kahrizi D, Bahrami G, Mohammadi-Motlagh HR. Study of gene expression and steviol glycosides accumulation in *Stevia rebaudiana* Bertoni under various mannitol concentrations. *Mol Biol Rep* 2019; 46(1): 7-16.

37. Eryugur N, Ucar E, Akpulat HA, Shahsavari K, Safavi SM, Kahrizi D. In vitro antioxidant assessment, screening of enzyme inhibi-

tory activities of methanol and water extracts and gene expression in *Hypericum lydium*. *Mol Biol Rep* 2019; 46(2): 2121-9.

38. Esmaeili F, Ghaheri M, Kahrizi D, Mansouri M, Safavi SM, Ghorbani T, Mohammadi S, Rahmanian E, Vaziri S. Effects of various glutamine concentrations on gene expression and steviol glycosides accumulation in *Stevia rebaudiana* Bertoni. *Cell Mol Biol*, 64(2): 1-5.

39. Akbari F, Arminian A, Kahrizi D, Fazeli A, Ghaheri M. Effect of nitrogen sources on gene expression of *Stevia rebaudiana* (Bertoni) under in vitro conditions. *Cell Mol Biol*, 64(2): 11-16.

40. Ghaheri M, Adibrad E, Safavi SM, Kahrizi D, Soroush A, Mohammadi S, Ghorbani T, Sabzevari A, Ansarypour Z, Rahmanian E. Effects of life cycle and leaves location on gene expression and glycoside biosynthesis pathway in *Stevia rebaudiana* Bertoni. *Cell Mol Biol*, 64(2): 17-22.

41. Kahrizi D, Ghaheri M, Yari Z, Yari K, Bahraminejad S. Investigation of different concentrations of MS media effects on gene expression and steviol glycosides accumulation in *Stevia rebaudiana* Bertoni. *Cell Mol Biol*, 64(2): 23-27.

42. Hashempoor S, Ghaheri M, Kahrizi D, Kazemi N, Mohammadi S, Safavi SM, Ghorbani T, Rahmanian E, Heshmatpanaah M. Effects of different concentrations of mannitol on gene expression in *Stevia rebaudiana* Bertoni. *Cell Mol Biol*, 64(2): 28-31.

43. Akbarabadi A, Ismaili A, Kahrizi D, Firouzabadi FN. Validation of expression stability of reference genes in response to herbicide stress in wild oat (*Avena ludoviciana*). *Cell Mol Biol* 2018; 64(4): 113-118.

44. Ghorbani T, Kahrizi D, Saeidi M, Arji I. Effect of sucrose concentrations on *Stevia rebaudiana* Bertoni tissue culture and gene expression. *Cell Mol Biol*, 63(11): 32-36.

45. Kahrizi D, Ghari SM, Ghaheri M, Fallah F, Ghorbani T, Beheshti AA, Kazemi E, Ansarypour Z. Effect of KH<sub>2</sub>PO<sub>4</sub> on gene expression, morphological and biochemical characteristics of *stevia rebaudiana* Bertoni under in vitro conditions. *Cell Mol Biol*, 63(7): 107-111.

46. Fallah F, Nokhasi F, Ghaheri M, Kahrizi D, Beheshti Ale AA, Ghorbani T, Kazemi E, Ansarypour Z. Effect of salinity on gene expression, morphological and biochemical characteristics of *Stevia rebaudiana* Bertoni under in vitro conditions. *Cell Mol Biol* 2017; 63(7): 102-106.

47. Fatemi SS, Yari K, Tavallaee M, Kahrizi D. Expression of recombinant Hc domain of *Clostridium botulinum* neurotoxin A in *E. coli* and its purification as a vaccine candidate against botulism. *Sci J Kurdistan Univ Med Sci* 2011 Aug 10;16(2):36-44.

48. Motamedi MJ, Shahsavandi S, Amani J, Kazemi R, Takrim S, Jafari M, Salmanian AH. Immunogenicity of the Multi-Epitopic Recombinant Glycoproteins of Newcastle Disease Virus: Implications for the Serodiagnosis Applications. *Iran J Biotechnol.* 2018; 16(4): e1749.

49. Ali S, Ahmad Nasir I, Rafiq M, Javed Butt S, Ihsan F, Qayyum Rao A, Husnain T. Sugarcane Mosaic Virus-Based Gene Silencing in *Nicotiana benthamiana*. *Iran J Biotechnol.* 2017; 15(4): 260-267.

50. Chen T, Zou X, Xu Z, Wang Y, Wang P, Peng H, Liu D, Lin J, Luo R, Wang Y, Chen Q, Chen D, Cai M, Li M. Molecular Characterization of the Epstein-Barr Virus BGLF2 Gene, its Expression, and Subcellular Localization. *Iran J Biotechnol.* May 2018; 16(2): e1610.

51. Rao X, Sun J. Development of SYBR Green I Based Real-Time RT-PCR Assay for Specific Detection of Watermelon silver mottle Virus. *Iran J Biotechnol.* Sep 2015; 13(3): 20-24.

52. Savardashtaki A, Sharifi Z, Hamzehlou S, Farajollahi MM. Analysis of Immunoreactivity of Heterologously Expressed Non-structural Protein 4B (NS4B) from Hepatitis C Virus (HCV) Genotype 1a. *Iran J Biotechnol.* Dec 2015; 13(4): 32-37.

53. Shafaghi M, Maktoobian S, Rasouli R, Howaizi N, Ofoghi H, Ehsani P. Transient Expression of Biologically Active Anti-rabies Virus Monoclonal Antibody in Tobacco Leaves. *Iran J Biotechnol.* Apr 2018; 16(1): e1774.
54. Shibaie N, Majidi J, Bashir NS, Karkhaneh A, Razavi K. Production and Partial Purification of the Grapevine Fanleaf Virus Coat Protein 42 Polyclonal Antibody Against Inclusion Body Expressed in *Escherichia coli*. *Iran J Biotechnol.* Dec 2018; 16(4): e1525.
55. Singh A, Permar V, Basavaraj, Tomar BS, Praveen S. Effect of Temperature on Symptoms Expression and Viral RNA Accumulation in Groundnut Bud Necrosis Virus Infected *Vigna unguiculata*. *Iran J Biotechnol.* Aug 2018; 16(3): e1846.
56. Sokhandan Bashir N, Poorsmaile M, Hajizadeh M. Heterologous Expression of Potato Virus Y Coat Protein, Isolate Pot187. *Iran J Biotechnol.* Dec 2015; 13(4): 48-52.
57. Vafaei SH, Mahmoodi M. Presence of Recombinant Strain of Cucurbit Aphid Borne Yellow Virus in Iran. *Iran J Biotechnol.* 2017; 15(4): 289-295.
58. Venkataravanappa V, Reddy CNL, Shankarappa KS, Reddy MK. Association of Tomato Leaf Curl New Delhi Virus, Betasatellite, and Alphasatellite with Mosaic Disease of Spine Gourd (*Momordica dioica* Roxb. Willd) in India. *Iran J Biotechnol.* Jan 2019; 17(1): e2134.
59. Yan Y, Jiang W, Liu J, Xu W, Qian H. Expression of Recombinant Phosphodiesterases 3A and 3B Using Baculovirus Expression System. *Iran J Biotechnol.* Dec 2016; 14(4): 236-242.
60. Zhang J, Jin Z, Sun T, Jiang Y, Han Q, Song Y, Chen Q, Xia X. Prokaryotic Expression, Purification, and Polyclonal Antibody Production of a Truncated Recombinant Rabies Virus L Protein. *Iran J Biotechnol.* Jun 2015; 13(2): 18-24.
61. Zou X, Xu Z, Wang Y, Ou X, Li Y, Liu D, Gan W, Lu M, Chen Q, Peng H, Hong G, Lin J, Li M, Cai M. Expression, Purification, and Antiserum Production of the Truncated UL31 Protein of Herpes Simplex Virus 1. *Iran J Biotechnol.* Jan 2019; 17(1): e1609.